Transactions in GIS

Guest Editorial

Virtual Environments Begin to Embrace Process-based Geographic Analysis

Virtual environments are computer-based digital spaces that we can observe, participate in, and experience in person. Virtual environments were initially viewed as "mirror worlds" (Gerlenter 2004), although they are being extended as this virtuality broadens to include scenarios that cannot be found in real life. Employing virtual environments as a tool to facilitate scientific exploration is not a new story. For example, virtual worlds, such as Second Life (SL) and World of Warcraft (WoW), have demonstrated the potential for studying social, behavioral and economic issues (Bambridge 2007; DeMers et al. 2013; Messinger et al. 2009), while a wide array of virtual laboratories are being developed to advance research in conventional domains, such as those dealing with chemical and biological problems (Bettenworth and Abbott 2000; Bohil et al. 2011; Yaron et al. 2010).

Similar examples have been reported in the domain of geography. Several three-dimensional geospatially referenced virtual environments (GRVEs) have been developed at different scales to include virtual objects (e.g. terrain, landmarks, and buildings) that correspond to real world representations. With these 'windows' on the real world, users can observe and move in abstract representations of the real places in which they live and work or in unfamiliar places that they have not experienced hitherto. This approach provides a valuable strategy for obtaining images of real geographic scenes or future scenarios, thus facilitating direct perception. However, this strategy does not yet allow for rigorous geographic analysis because a mismatch exists between actual environments and the limited functions that GRVEs can provide (MacEachren and Kraak 2001).

Later developments have revealed a greater potential for using GRVEs as experimental laboratories for geospatial analysis because more theory and analytic functions have been incorporated into their structure. In enhanced GRVEs, spatial databases and indices have been employed so that users can easily query physical geographic objects (such as mountains, lakes, and buildings) and their attached attributes. Three-dimensional spatial analysis methods provide appropriate representations for computing the parameters of navigation that are associated with movement in both real and abstract space as well as various features, such as the average duration over which a building receives sunlight each day and the extent to which land-use activities are patronized by those working and/or living in such environments. Moreover, with the development of three-dimensional physical data models, geographers can perform cross-sectional analyses of virtual geological and geographic structures from multiple directions across many sub-layers. All these achievements focus on the question 'what role will virtual systems play in learning about the world?' (Cutter et al. 2002).

A key driver for enhancing assistive tools must stem from the target itself. Recently, an important research priority in geographic analysis has emerged as researchers shift from identifying static snapshots to exploring dynamic phenomena (Goodchild 2009, 2010). Dynamic phenomena, unlike static shapes, forms, and patterns, include a series of physical or human geographic processes that change across spatial and temporal dimensions. Manipulating, interpreting, and storing information for those changing processes are critical skills and are also

Acknowledgements: We thank Professor Michael F Goodchild for his valuable suggestions. The paper was supported by the National Key Basic Research Program of China (2015CB954103) and PAPD.

important for geographic understanding and analyses (Mark et al. 1999). This change in approach covers many aspects, ranging from theories to technologies, and provides new opportunities and challenges for related assistive tools and interfaces, including GRVEs.

New ideas are emerging with the aim of introducing at least three categories of geographic processes that are becoming embedded in GRVEs: historical geographic processes recorded in traditional documents, contemporary geographic processes that currently occur in nature, and experimental geographic processes that are still being explored and discovered.

Historical geographic processes pertain to changes that have occurred in the past, for example, the evolution of a city or the migratory process associated with a family. By adding historical geographic processes to a virtual environment, their historiography can be improved and participants can easily begin to compare temporal changes within a geographic context that embrace culture, the arts and humanities. However, most geographic information related to such processes is recorded in traditional paper maps, paintings, and unstructured texts, which cannot be directly used for GRVEs. Some attempts have been made to build four-dimensional representations (where time constitutes the fourth dimension), such as historical evolutions of virtual cities, e.g. virtual Kyoto (Yasumoto et al. 2012), to draw four-dimensional migration paths of Chinese families (Chen et al. 2013), and so on; but accuracy and precision remain problems for further study.

The perceptual psychologist Gibson argued that a good laboratory must resemble life itself (Gibson 1986). Following Gibson, arguably, a suitable virtual environment should be as lifelike as possible (Loomis et al. 1999; Tarr and Warren 2002). One's surroundings that embody physical processes, such as blowing wind, pouring rain, moving cars, and animals in motion, inform our daily life in a dynamic world. Introducing this information into GRVEs will enhance the user's perception of the environment and will enable better feedback and reactions from users (Lin et al. 2013b).

In this respect, accessing up-to-date geographic process data is a major quest. Currently, such process data can be acquired at increasingly faster rates. Environmental data (e.g. mineral and geothermal resources, air and water quality, humidity, and temperature) can be collected by ubiquitous sensors, while social data (e.g. travel behavior, energy use, and spending patterns) can be gathered using various forms of crowdsourcing, which can supplement remote sensing observations. With timely positioning, these data can be an important component of GRVEs, providing facts and connections for the users. For example, in Figure 1, we show how such a virtual environment can be created by importing into a 3-D environment real-time streamed data from travel using smart cards, demonstrating how users of a particular category, i.e. those with a disabled free pass for travel, move around a complex system composed of buildings and subway lines that provide a virtual representation of the real world (Batty 2013).

Apart from this type of visualization, another important goal for geographically referenced virtual environments is their use as an experimental workspace for exploring real geographic phenomena (Bainbridge 2007; Subrahmanian and Dickerson 2009). Focusing on geographic research, computer-aided geographic experiments are quickly becoming a common strategy for simulating and predicting dynamic phenomena, where process models are often employed as basic tools. Geographic process models equip GRVEs with realistic mechanisms that allow users to perform exploratory analysis based on experiences and interactions with visual predictions from geographic scenarios that not yet well understood (Lin et al. 2013a). During such experiments researchers can adjust model-related parameters that pertain, for example, to the objects that comprise a city associated with air pollution (Xu et al. 2013) or the speed and flexibility of people as agent-based avatars interacting with a scene and exploring ways that they might move in such environment (Torrens 2012). Users can then compare different experimental results using straightforward visual perceptions.



Figure 1 A virtual, geographically referenced environment in which real-time data on movements illustrate flows of a particular class of traveler on London's subway systems (courtesy of Gareth Simons, CASA, University College London: http://vimeo.com/96998519).

Moreover, as a foundation for collective modeling and simulation, technologies related to model management, reuse, and coupling are being explored (Kelly et al. 2013; Wen et al. 2013). By incorporating such technologies, GRVEs can be improved as tools for researchers to share knowledge and gain a more universal understanding of geographic processes (Lin et al. 2013a; Bilke et al. 2014). A lower threshold can be used for comprehensive experiments in a collective mode, such as those being developed for flood-routing (Lin et al. 2010) or air pollution analysis (Xu et al. 2011).

Research has shown that the most effective way to understand an environment is from direct experience (Cutter et al. 2002; Tuan 2001). Massive geographic datasets are becoming increasingly available for representing such environments, while the causality for expressing various processes is becoming better understood when combined with such information and integrated in future analysis. These efforts will provide more realistic and perceptible GRVEs, a holograph for geography so-to-speak, that allow users to explore inaccessible past or future environments or distant present environments, not only through their static objects but through processes that mirror their real dynamics. Ongoing research is challenging because it involves many technologies in the domains of geography, virtual reality and computer science that must be synthesized from both physical and human perspectives. However, these challenges should not obscure the fact that these new research directions will lead to a new synthesis of the virtual and real worlds. Our speculation is that embedding geographic processes into virtual environments holds extraordinary potential for future geographic analysis, which extends well beyond the current state of the art in geographic information systems and geographic information science.

Hui Lin

Institute of Space and Earth Information Science The Chinese University of Hong Kong; Shenzhen Research Institute The Chinese University of Hong Kong; Department of Geography and Resource Management, The Chinese University of Hong Kong

Michael Batty Centre for Advanced Spatial Analysis University College London

Sven E Jørgensen Faculty of Health, Institute A, Section of Environmental Chemistry University of Copenhagen

> Bojie Fu State Key Laboratory of Urban and Regional Ecology Research Center for Eco-environmental Sciences Chinese Academy of Sciences

> > Milan Konecny Laboratory on Geoinformatics and Cartography Masaryk University

> > > Alexey Voinov Faculty of Geo-Information Science and Earth Observation (ITC) University of Twente

Paul Torrens Center for Geospatial Information Science University of Maryland

Guonian Lu Key Laboratory of Virtual Geographic Environment, Ministry of Education of PRC, Nanjing Normal University

A-Xing Zhu Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing Normal University; Department of Geography, University of Wisconsin–Madison

> John P Wilson Spatial Sciences Institute University of Southern California

> > Jianya Gong LIESMARS Wuhan University

Olaf Kolditz Department Environmental Informatics Technische Universität Dresden

© 2015 John Wiley & Sons Ltd

Temenoujka Bandrova Laboratory on Cartography University of Architecture, Civil Engineering and Geodesy

Min Chen

Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing Normal University; State Key Laboratory Cultivation Base of Geographical Environment Evolution (Jiangsu Province), Nanjing Normal University Address for correspondence: chenmin0902@163.com

References

Bainbridge W S 2007 The scientific research potential of virtual worlds. Science 317: 472-76

- Batty M 2013 The New Science of Cities. Cambridge, MA, MIT Press
- Bettenworth V and Abbott A 2000 Virtual laboratory will recreate classic biology experiments. Nature 408: 761
- Bilke L, Fischer T, Helbig C, Krawczyk C, Nagel T, Naumov D, Paulick S, Rink K, Sachse A, Schelenz S, Walther M, Watanabe N, Zehner B, Ziesch J, and Kolditz O 2014 VISLAB: Laboratory for scientific visualization. *Environmental Earth Sciences* 72: 3881–99
- Bohil C J, Alicea B, and Biocca F A 2011 Virtual reality in neuroscience research and therapy. Nature Reviews Neuroscience 12: 752-62
- Chen M, He L, Lin H, Zhang C, and Hu M 2013 Real-geographic-scenario-based virtual social environments: Integrating geography with social research. *Environment and Planning B* 40: 1103–21
- Cutter S L, Golledge R, and Graf W L 2002 The big questions in geography. Professional Geographer 54: 305-17
- DeMers M, Klimaszewski-Patterson A, Richman R, Ahearn S, Plewe B, and Skupin A 2013 Toward an immersive 3D virtual BoK Exploratorium: A proof of concept. *Transactions GIS* 17: 335–52
- Gerlernter D 2004 Mirror Worlds: or the Day Software Puts the Universe in a Shoebox ... How It Will Happen and What It Will Mean. New York, Oxford University Press
- Gibson J J 1986 The Ecological Approach to Visual Perception (Third Edition). Boston, MA, Houghton Mifflin

Goodchild M F 2009 Geographic information systems and science: Today and tomorrow. Annals of GIS 15: 3-9

- Goodchild M F 2010 Twenty years of progress: GIScience in 2010. Journal of Spatial Information Science 1: 3-20
- Kelly (Letcher) R A, Jakeman A J, Barreteau O, Borsuk M E, Elsawah S, and Hamilton S H 2013 Selecting among five common modelling approaches for integrated environmental assessment and management. *Environmen*tal Modeling and Software 47: 159–81
- Lin H, Chen M, and Lu G 2013a Virtual geographic environment: A workspace for computer-aided geographic experiments. *Annals of the Association American Geographers* 103: 465–82
- Lin H, Chen M, Lu G, Qing Zhu Q, Gong J, You X, Wen Y, Xu B, Hu M 2013b Virtual geographic environments (VGEs): A new generation of geographic analysis tool. *Earth-Science Reviews* 126: 74–84
- Lin H, Zhu J, Gong J, Xu B, and Qi H 2010 A grid-based collaborative virtual geographic environment for the planning of silt dam systems. *International Journal of Geographical Information Science* 24: 607–21
- Loomis J M, Blascovich J J, and Beall A C 1999 Immersive virtual environment technology as a basic research tool in psychology. *Behavior Research Methods, Instruments, and Computers* 31: 557–64
- MacEachren A M and Kraak M 2011 Research challenges in geovisualization. Cartography and Geographic Information Science 28: 3–12
- Mark D M, Freksa C, Hirtle S C, Lloyd R, and Tversky B 1999 Cognitive models of geographical space. International Journal of Geographical Information Science 13: 747–74
- Messinger P R, Stroulia E, Lyons K, Bone M, Niu R H, Smirnov K, and Perelgut S 2009 Virtual worlds past, present, and future: New directions in social computing. *Decis. Support Systems* 47: 204–28
- Subrahmanian V S and Dickerson J 2009 What can virtual worlds and games do for national security? *Science* 326: 1201–02
- Tarr M J and Warren W H 2002 Virtual reality in behavioral neuroscience and beyond. *Nature Neuroscience* 5 (Suppl.): 1089–92
- Torrens P M 2012 Moving agent pedestrians through space and time. Annals of the Association of American Geographers 102: 35-66

498 *Lin et al.*

Tuan Y-F 2001 Space and Place: the Perspective of Experience. Minneapolis, MN, University of Minnesota Press

- Wen Y, Chen M, Lu G, and Lin H 2013 Prototyping an open environment for sharing geographical analysis models on cloud computing platform. *International Journal of Digital Earth* 6: 356–82
- Xu B, Lin H, Chiu L, Hu Y, Zhu J, Hu M, and Cui W 2011 Collaborative virtual geographic environments: a case study of air pollution simulation. *Information Sciences* 181: 2231–46
- Xu B, Lin H, Gong J, Tang S, Hu Y, Nasser I A, and Jing T 2013 Integration of a computational grid and virtual geographic environment to facilitate air pollution simulation. *Computers and Geosciences* 54: 184–95
- Yaron D, Karabinos M, Lange D, Greeno J G, and Leinhardt G 2010 The ChemCollective: Virtual labs for introductory chemistry courses. *Science* 328: 584–85
- Yasumoto S, Jones A, Yano K, and Nakaya T 2012 Virtual city models for assessing environmental equity of access to sunlight: A case study of Kyoto, Japan. *International Journal of Geographical Information Science* 26: 1–13